

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/073686 A2

- (51) International Patent Classification⁷: **A61K 9/14** (74) Agent: **BRAINARD, Charles, R.**; Kenyon & Kenyon, One Broadway, New York, NY 10004-1050 (US).
- (21) International Application Number: **PCT/US2004/004684** (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 17 February 2004 (17.02.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/449,246 20 February 2003 (20.02.2003) US
- (71) Applicant (for all designated States except BB, US): **TEVA PHARMACEUTICAL INDUSTRIES LTD.** [IL/IL]; 5 Basel Street, P.O. Box 3190, 49131 Petah-Tiqva (IL).
- (71) Applicant (for BB only): **TEVA PHARMACEUTICALS USA, INC.** [US/US]; 1090 Horsham Road, P.O. Box 1090, North Wales, PA 19454-1090 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **FLASH-NER-BARAK, Moshe** [IL/IL]; Hefetz Mordechai 15, Petach Tikav (IL). **LERNER, Itzhak, E.** [IL/IL]; Wolfson 32, Petach Tiqva (IL). **ROSENBERGER, Vered** [IL/IL]; 3 Miss Landau Street, Jerusalem 96410 (IL). **MOLDAVSKI, Naomi** [IL/IL]; Moshav Dekel D.N., HaNegev 85492 (IL).
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MENTHOL SOLUTIONS OF DRUGS

(57) Abstract: The present invention relates to compositions comprising solutions of drugs in menthol, especially drugs that are poorly soluble in water, and to methods for making such compositions.

WO 2004/073686 A2

MENTHOL SOLUTIONS OF DRUGS

Field of the Invention

The present invention encompasses compositions comprising solutions or solid solutions of drugs in menthol, especially drugs that are poorly soluble in water, and to methods for making such compositions.

Background of the Invention

Several clinically important drugs have limited oral bioavailability and high interpatient variability, resulting in difficulty in obtaining optimum treatment regimens for their use. Reasons for such limited oral bioavailability may include poor solubility in water or biological fluids, poor membrane permeability, efficient MDR (multiple drug resistance) pumps, and/or destructive metabolism in the intestine or the liver. Such metabolic destruction may be by the family of cytochrome P450 enzymes that oxidatively destroy many drugs (*e.g.* CYP3A4) or by glucuronidation enzymes that help the body eliminate the glucuronide derivatives of the drug in the urine or by excretion in the bile to the feces. High interpatient variability is often associated with the genetic variability of metabolic pathways in humans as well as the genetic variation in the expression of the P-glycoprotein MDR pumps.

Drugs with limited oral bioavailability include cyclosporines. Cyclosporines are a very important family of drugs which are used for the avoidance of organ rejection after organ transplant. Cyclosporines, however, suffer from erratic absorption caused by most of the factors mentioned above. See, A. Lindholm, "Factors Influencing the Pharmacokinetics of Cyclosporine in Man," *Therapeutic Drug Monitoring*, 13 (6), 465-477 (1991). Cyclosporines are insoluble in water, are expelled from cells of the intestine by P-glycoprotein efflux pumps, and are heavily metabolized both in the intestine and in the liver by cytochrome P-450 enzymes. Ducharme, *et al.*, "Disposition of Intravenous and Oral Cyclosporine after Administration with Grapefruit Juice," *Clinical Pharmacology and Therapeutics*, 57(5), 485-491 (1995); and Wu, *et al.*, "Differentiation of Absorption and First - Pass Gut and Hepatic Metabolism in Humans: studies with Cyclosporine," *Clinical Pharmacology and Therapeutics*, 58(5), 449-497 (1995). Since the therapeutic window for cyclosporines is not very wide and the toxic effects of overdose are pronounced dosing with this drug type has traditionally been difficult. See *e.g.*, PHYSICIAN'S DESK REFERENCE, pp. 2310-2313 (57th Ed. 2003).

Cyclosporines were originally formulated in oil-based formulations so as to dissolve the drug. Oil and water do not mix very well, thus adding to the variability of the bioavailability of the product. The use of cyclosporines in microemulsion formulations has somewhat improved this situation, however, the efflux pump and oxidative metabolism issues remain essentially as problematical as before. To address the poor bioavailability of cyclosporines and other drugs, Benet and co-workers described dosing the drug either after, with, or mixed with essential oils or essential oil components such as menthol and carvone among others. See, Benet, *et al.*, U.S. Patent Nos.: 5,665,386; 5,716,928; 6,121,234; 6,004,927; and 6,028,054. Benet showed, using in vitro tests, an inhibition of metabolism of cyclosporine and other drugs and a concomitant improvement in bioavailability. The insolubility of the drug in water and the incompatibility of oil based formulations with the aqueous environment of the human gut were still present.

The pharmacokinetics of cyclosporine have been studied using the classical oil-based formulation and the improved microemulsion formulation along with metabolic inhibitors such as ketoconazole. Aklaghi, *et al.*, "Pharmacokinetics of Cyclosporine in Heart Transplant Recipients Receiving Metabolic Inhibitors," *The Journal of Heart and Lung Transplantation*, 20 (4), 431-438 (2001). Ketoconazole inhibits CYP3 metabolism as well as P-glycoprotein efflux pumps. The microemulsion formula gave improved bioavailability and somewhat of an improvement in variability when tested without ketoconazole. Treatment with ketoconazole greatly improved the bioavailability of the cyclosporine but not the variability. When pretreating with ketoconazole, the micro emulsion formulation was no better than the oil based formulation. While concurrent treatment with ketoconazole is practiced in certain medical centers to improve cyclosporine bioavailability, in general, the medical community is against giving potent drugs with serious toxic side effects as an adjuvant for another drug when there is no medical need for its administration. Ketoconazole is a potent anti-fungal which is known to exhibit side effects. The need for a safe alternative that will both raise the bioavailability of cyclosporines and lower the inter-patient variability is still present.

The statin drugs, which are used to treat high cholesterol levels, have become some of the most widely used drugs in the world. The family of statin drugs suffers from poor oral bioavailability. This poor oral bioavailability is believed to be caused, to a great extent, by high first pass metabolism. Simvastatin, one of the most widely used drugs in this class, is a prodrug of its active metabolite. However, only about 5% of the dose is

available as the active metabolite in the blood due to hepatic first pass metabolism.

MARTINDALE: THE COMPLETE DRUG REFERENCE, pp. 969-970 (33rd Ed., 2002). The statins have serious toxic side effects in terms of muscle disorders, rhabdomyolysis being one of the more serious side effects. *Id.* As with cyclosporines, the inter-patient
5 variability makes it difficult for the doctor to tailor proper dosage for the patient so as to give effective cholesterol reduction without toxic adverse events. Simvastatin has been administered with grapefruit juice or with a capsule of peppermint oil. Peppermint oil, which is known to inhibit CYP3A4, raised bioavailability 60% while the grapefruit juice, which is known to inhibit both CYP3A4 and the P-glycoprotein efflux pump, raised
10 bioavailability 300%. Wacher, *et al.*, "Peppermint Oil Increases the Bioavailability of Felodipine and Simvastatin," *Clinical Pharmacology and Therapeutics*, 71(2), P67 Abstract TPII-95.

Paclitaxel is an important antineoplastic agent that is administered by intravenous injection. Paclitaxel suffers from very poor solubility in water. The insolubility hampers
15 i.v. dosing, causing a need for special formulations which may have non-trivial toxicity profiles. PHYSICIAN'S DESK REFERENCE, pp. 1129-1138 (56th Ed. 2002). The insolubility of paclitaxel also hampers use in oral dosing. This problem, however, is minor in comparison to the effects of the P-glycoprotein efflux pump in the intestine. Paclitaxel has been successfully dosed orally by co-administering it with efficient
20 inhibitors of the P-glycoprotein pump such as cyclosporines. See, Malingre, *et al.*, "The Effect of Different Doses of Cyclosporin A on the Systemic Exposure of Orally Administered Paclitaxel," *Anti-Cancer Drugs*, 12, 351-358 (2001); Malingre, *et al.*, "A Phase I and Pharmacokinetic Study of Bi-Daily Dosing of Oral Paclitaxel in Combination with Cyclosporin A," *Cancer Chemother Pharmacol.*, 47, 347-354 (2001); and Broder, *et al.*, U.S. Patent Nos.: 5,968,972; and 6,395,770. Cyclosporines, however, are much too
25 potent a drug type to be used as an adjuvant for the enhanced availability of another drug, even one as important as paclitaxel. Another method of orally dosing paclitaxel is clearly needed.

Many drugs have glucuronidation as their main metabolic pathway of elimination.
30 GOODMAN AND GILMAN'S: THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, p. 13 (9th ed., 1996); and De Wilt, *et al.*, "Glucuronidation in Humans. Pharmacogenetic and Developmental Aspects," *Clinical Pharmacokinetics*, 36(6), 439-452 (1999). Recent evidence shows that this pathway may be important in the metabolism of simvastatin along with the mechanisms described above. Prueksaritanont, *et al.*, "Glucuronidation of

Statins in Animals and Humans: A Novel Mechanism of Statin Lactonization,” *Drug Metabolism and Disposition*, 30, 505-512 (2002). The present invention overcomes many of the existing limitations within the prior art by providing novel formulations.

Summary of the Invention

5 One embodiment of the invention encompasses compositions for increasing the oral bioavailability of a drug comprising at least one poorly bioavailable drug dissolved in an effective amount of menthol. The poorly bioavailable drug may be at least one drug with low aqueous solubility, a drug metabolized by cytochrome P450, a drug expelled from cells by the P-glycoprotein pump, or a drug metabolized via glucuronidation. A
10 drug with low aqueous solubility is a drug having a water solubility of less than about 20 mg/ per milliliter of water.

Another embodiment of the invention encompasses compositions wherein the poorly bioavailable drugs include, but are not limited to, at least one of cyclosporine, atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pravastatin, simvastatin,
15 paclitaxel, fenofibrate, itraconazole, bromocriptine, carbamazepine, diazepam, paclitaxel, etoposide, camptothecin, danazole, progesterone, nitrofurantoin, estradiol, estrone, oxfendazole, proquazone, ketoprofen, nifedipine, verapamil, or glyburide. Preferably, the drug includes cyclosporine, atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pravastatin, simvastatin, or paclitxel. More preferably, the drug is simvastatin.

20 Yet another embodiment of the invention encompasses methods for improving the bioavailability of a drug comprising dissolving the drug in an effective amount of menthol. Another method of the invention encompasses methods for improving the oral bioavailability of a drug comprising dissolving at least one poorly bioavailable drug in an effective amount of menthol. The method may further comprise administering the
25 composition to a mammal. In one embodiment, the amount of menthol sufficient to increase the drug's bioavailability may be from about 20% to about 99% by weight, preferably, the menthol may be present in an amount of about 60% to about 95% by weight of the composition. Alternatively, the amount of menthol may be sufficient to increase the oral bioavailability of the drug by an increase of about 10% or more in the
30 average area under the blood or plasma concentration versus time curve (AUC) when compared to the average AUC for a non-menthol containing composition of the drug.

Yet another embodiment of the invention encompasses methods for reducing the variability of the bioavailability of a drug comprising dissolving at least one poorly

bioavailable drug in an effective amount of menthol. The method may further comprise administering the composition to a mammal. In one embodiment of the method, the amount of menthol may be sufficient to decrease the variability in the drug's bioavailability by about 10% or more of the relative standard deviation (CV%) of the area under the blood or plasma concentration versus time curve (AUC) when compared to the AUC of a non-menthol containing formulation of the drug.

Another embodiment of the invention encompasses methods for increasing the extent of time that a drug provides a therapeutically significant concentration in blood or plasma comprising dissolving at least one poorly bioavailable drug in an effective amount of menthol. In one embodiment, the amount of menthol may be sufficient to extend the time that the drug provides a therapeutically significant concentration in blood or plasma by one hour or more.

Detailed Description of the Invention

The invention comprises formulations of drugs with low bioavailability and menthol. As used herein, the term "poor bioavailability" or "poorly bioavailable" refers to a drug that has an oral bioavailability in its active form, whether it be the drug as dosed or an active metabolite thereof, of less than 30%.

Not to be limited by theory, it is believed that the compositions of the invention operate, in part, by providing a composition where the poorly bioavailable drugs are combined with compounds that aid solubility and/or compounds that compete with the poorly bioavailable drug in the biodegradable pathway which degrades the poorly bioavailable drugs. The delivery of the poorly bioavailable drugs is improved by using materials that are generally recognized as safe and without the use of potent drugs to establish an efficient competition within the biodegradable pathway. Thus, the non-active compound would be metabolized prior to the active drug. In particular, our studies found that formulating poorly bioavailable drugs as a solution or a solid solution in menthol improved delivery as compared to dosing the drug alone, dosing the drug after a menthol dose or a menthol containing dose (e.g. peppermint oil), or dosing the drug along with a dose of menthol. The compositions of the invention allow for the use of lower doses of drugs that provide the same systemic concentrations of drugs as the currently supplied doses that undergo extensive presystemic metabolism and degradation. Also, the compositions of the invention reduce interpatient variability caused by the inherently differing metabolic profiles between subjects.

Menthol, chemically known as (1 α ,2 β ,5 α)-5-methyl-2-(1-methylethyl)-cyclohexanol, is partially soluble in water. Because menthol has a low melting point, *i.e.*, about 41°C to 43°C, compositions of menthol and drugs dissolved within the menthol have melting points close to body temperature. This property allows menthol to act as an efficient solvent for many drugs. We have found that menthol is a superior solvent for poorly water soluble drugs as compared to oil based drug formulations, because, in part, the drugs are more available to the aqueous medium of the gastro-intestinal tract as compared to oil based formulations. Although, menthol has been known to act as a skin absorption enhancer, it is believed that menthol may also improve gastro-intestinal drug absorption as well.

The invention advantageously uses menthol in close proximity with poorly bioavailable drugs to deter drug biodegradation in a kinetically competitive environment. In other words, menthol may be used to inhibit biological degradation pathways which metabolize the active drug and/or kinetically compete with the drug at the biologically active degradation site. For example, menthol inhibits CYP3A4 metabolism and the P-glycoprotein pump, thus, menthol in close proximity to and in intimate contact with the poorly bioavailable drug greatly enhances the bioavailability of the drug as the drug does not undergo degradation. Also, menthol which has been shown to be metabolized to a glucuronide derivative, can serve as a sacrificial molecule wherein menthol is degraded prior to the drug, thus delaying drug degradation and extending drug bioavailability. In other words, menthol is potentially capable of competing with a drug as a decoy for glucuronidation, thereby leaving less of the drug metabolized and yielding an overall increase in the drug bioavailability.

The present invention encompasses pharmaceutical compositions for improving the bioavailability of a drug comprising at least one drug dissolved in an effective amount of menthol. In particular, the invention encompasses pharmaceutical compositions for improving the bioavailability of a drug comprising at least one poorly bioavailable drug dissolved in an effective amount of menthol. As used herein, the term "improving bioavailability" refers to the increase in concentration of a drug as compared to the concentration of the drug without menthol. In other words, drug bioavailability is proportional to, and is typically measured by, the total area under the curve (AUC) of the concentration of the drug found in blood or plasma versus time when measured in a pharmacokinetic trial in a human or an animal. The AUC may be expressed as AUC_t,

i.e. the area under the curve to the last measured time point, or AUC_t , *i.e.* the area under the curve extrapolated to infinite time. The improvement in bioavailability is measured by the percent increase in the average AUC of the subjects in the trial when dosing the drug dissolved in menthol as compared to the average AUC of the same subjects obtained by standard dosing of the drug. Alternately, the AUC ratio of the test formulation (AUC_f) to the AUC of the reference formulation (AUC_r) may be calculated on a per subject basis and then averaged. A percent of the average ratio (AUC_f/AUC_r) above 100% is then the improvement in bioavailability. Typically, the improvement in the average AUC when dosing the drug dissolved in menthol as compared to the average AUC obtained by standard dosing of the drug is about 5%, and preferably, the improvement is about 10% or more in the bioavailability, which is considered significant.

The present invention further provides a pharmaceutical composition directed to improving the extent of time that a drug provides a therapeutically significant concentration in blood or plasma and/or reducing the drug bioavailability variability, wherein the drug is dissolved in menthol. As used herein, the term "improving the extent of time" refers to the increase in length of time that a drug provides a therapeutically significant concentration in blood or plasma. Preferably, the time a drug provides a therapeutically significant concentration in blood or plasma is extended by about one hour or more. As used herein, the term "drug bioavailability variability" is defined as the relative standard deviation, expressed as CV%, of the drug's AUC over the subjects tested. A highly variable drug is one with a CV% greater than 50%. An improvement of the CV% by 10 percent or more is considered significant. The present invention is particularly directed to a pharmaceutical composition comprising a solid or solid solution of a drug dissolved in an effective amount of menthol. The solid solution may include a compound or polymer that forms a dispersion with the drug.

The poor bioavailability of the drug may be due to several factors. Such factors include, but are not limited to, low aqueous solubility, metabolism by cytochrome P450, expulsion from cells by the P-glycoprotein pump, or metabolism via glucuronidation. Thus, the present invention encompasses compositions for increasing the bioavailability of drugs with low aqueous solubility, drugs metabolized by cytochrome P450, drugs expelled from cells by the P-glycoprotein pump, and/or drugs metabolized via glucuronidation. As used herein, the term "low aqueous solubility" refers to a drug that is considered to be poorly water-soluble, *i.e.*, the drug has a water solubility of less than about 20 mg/ per milliliter of water.

Any pharmacologically active substance or drug can be used in the practice of the present invention. Preferred drugs, however, include drugs having poor bioavailability. Examples of drugs having poor bioavailability include, but are not limited to, cyclosporine, statins, paclitaxel, fenofibrate, itraconazole, bromocriptine, carbamazepine, diazepam, paclitaxel, etoposide, camptothecin, danazole, progesterone, nitrofurantoin, estradiol, estrone, oxfendazole, proquazone, ketoprofen, nifedipine, verapamil, or glyburide. Statins include, but are not limited to, atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pravastatin, or simvastatin. Preferably, the drugs having poor availability include at least one of cyclosporine, statins, or paclitaxel. A more preferred statin is simvastatin. Other examples of drugs having poor bioavailability will be readily apparent to one of ordinary skill in the art.

The amount of drug in the composition of the invention should be sufficient to be therapeutically effective for the condition administered. One of ordinary skill in the art can easily determine with little or no experimentation the effective amount of drug.

Typically, the drug is present in an amount of about 5% to about 40% by weight of the composition, preferably, the drug is present in an amount of about 10%.

The amount of menthol in the composition of the invention should be sufficient to improve the bioavailability of the poorly bioavailable drug. Typically, the amount of improvement should be at least about 5% of the average AUC as compared to the average AUC of a non-menthol containing formulation and preferably, the improvement is about 15%. One of ordinary skill in the art can easily determine with little or no experimentation the effective amount of menthol. Typically, menthol is present in the composition in a amount of about 20% to about 99% by weight of the composition, and preferably, menthol is present in an amount of about 60% to about 95%. More preferably, menthol is present in the composition in an amount of about 80% to about 90% by weight.

The compositions of the invention may also encompass other excipients commonly used in drug manufacture including, but not limited to, binders, fillers, disintegrants, lubricants, colorants, carriers, and diluents.

Another embodiment of the invention encompasses methods of improving the bioavailability of a drug comprising dissolving the drug in an effective amount of menthol. In particular, the invention encompasses methods for improving the bioavailability of a drug comprising dissolving at least one drug with low aqueous solubility, drug capable of being metabolized by cytochrome P450, a drug capable of

being expelled from cells by the P-glycoprotein pump, or a drug capable of being metabolized via glucuronidation in an effective amount of menthol. Typically, the amount of improvement should be at least about 5% of the average AUC as compared to the average AUC of a non-menthol containing formulation and preferably about 15%, as explained above.

The invention encompasses methods for reducing the variability of the bioavailability of a drug comprising dissolving at least one drug with low aqueous solubility, a drug capable of being metabolized by cytochrome P450, a drug capable of being expelled from cells by the P-glycoprotein pump, or a drug capable of being metabolized via glucuronidation in an effective amount of menthol. As described above, drug variability is defined as the relative standard deviation, expressed as CV%, of the drug's AUC over the subjects tested. A highly variable drug is one with a CV% greater than 50%. Typically, the reduction is about 5% of the relative standard deviation (CV%) of the area under the blood or plasma concentration versus time curve (AUC) when compared to a non-menthol containing formulation average AUC, and preferably, the decrease in CV% is by about 10% or more, which is considered significant.

Another embodiment of the invention encompasses methods for increasing the extent of time that a drug provides a therapeutically significant concentration in blood or plasma comprising dissolving at least one poorly bioavailable drug in an effective amount of menthol. Typically, the extent of the bioavailability of a drug is increased by the administration of a composition comprising at least one drug and menthol, wherein the menthol is present in an amount sufficient to extend the time that the drug provides a therapeutically significant concentration by one hour or more.

The present invention encompasses unit dosage forms of the pharmaceutical composition comprising a unit dosage form of a drug dissolved in an effective amount of menthol. The compositions of the invention may be administered to a mammal. Preferably, the mammal is a human.

One embodiment encompasses the compositions of the invention be prepared into solid solution dosage forms. In particular, the compositions may be formulated into oral solid dosage forms such as capsules, tablets, or gelcaps. In particular, the pharmaceutical compositions can be made into unit dosage forms.

In one embodiment, the solid solution is formed on the surface of at least one pharmaceutical carrier particle. For example, a molten combination of drug and menthol can be applied to the surface of particles of one or more pharmaceutical carriers, and

allowed to cool to form the solid solution on the surface of the pharmaceutical carrier or carriers.

Having described the invention with reference to certain preferred embodiments, other embodiments will become apparent to one skilled in the art from consideration of the specification. The invention is further defined by reference to the following examples describing in detail the preparation of the composition and methods of use of the invention. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the scope of the invention.

Examples

Example 1

Cyclosporine (20 g) was heated in menthol (80 g) to 56°C while stirring until the cyclosporine dissolved yielding a clear solution. Microcrystalline cellulose (Avicel pH 102, 100 g) was added to the clear solution which was cooled to room temperature giving a solid solution of cyclosporine in menthol on the microcrystalline cellulose. The solid was milled using a Quadro Comil milling machine, with screens of 6350, 1575 and 813 microns sequentially used to produce a powder ready for filling into capsules.

Example 2

Simvastatin (20 g) was heated in menthol (200 g) to 60° C while stirring at 150 rpm in a jacketed reactor. The simvastatin dissolved in the menthol to give a clear solution. The solution was cooled to room temperature to a solid solution of simvastatin in menthol. The solid solution was milled using a Quadro Comil milling machine with a 1640 micron screen. The powder (200 mg) was filled into #0 capsules. The capsules were assayed for simvastatin content by dissolving a capsule in a pH 4 phosphate buffer containing acetonitrile (1:1). The simvastatin content was assayed on a C-18 column by HPLC and found to contain 20 mg of simvastatin per capsule. The release of simvastatin was measured in 450 ml of pH = 7 phosphate buffer containing 0.5% sodium lauryl sulfate (SLS) in water at 37°C and 50 rpm in an USP apparatus II dissolution system. The release was found to be greater than 75% at 30 minutes.

Example 3

Raloxifene HCl (60 mg, Evista, ELI LILLY®) was dosed to twelve healthy volunteers either alone or with a capsule containing 180 mg menthol in a crossover fashion with a two week washout between sessions. Blood samples were taken at 0, 0.5, 1, 2, 4, 6, 8, 10, 12, 16, 20, 24, 48, 72 and 96 hours and the content of raloxifene assayed.

5 The average C_{max} of the raloxifene dosed with menthol was 36% higher than the reference (320 pg/ml vs. 235 pg/ml), while the average area under the curve (AUC) was 8% higher when dosing with menthol (3041 vs. 12090 pg* hr/ml). Raloxifene is a long half-life drug ($t_{1/2}$ for the test was 26 hours and for the reference was 28 hours), while menthol has a short half-life. Without wishing to be bound by theory or mode of action,

10 it is believed that the main effect of menthol is seen in the first hours where it can effectively compete with the drug for glucuronization. An analysis of the AUC over the first six hours shows that the test AUC is 35% higher than the reference, mirroring the C_{max} result. Without wishing to be bound by theory or mode of action, it is believed that dosing with menthol can successfully compete with the metabolism of the drug, yielding

15 a better pharmacokinetic profile.

Example 4

An open-label study with randomized three-way crossover comparative pharmacokinetic study was conducted with 12 healthy fasted volunteers each receiving a

20 single dose of either: Reference-simvastatin (Simvastatin-Teva®, 20 mg) alone; Test 1-simvastatin (Simvastatin-Teva®, 20 mg) + menthol (180 mg capsule); or Test 2-simvastatin/menthol (10% simvastatin dissolved in menthol, 20 mg of simvastatin per capsule). A dose was administered to each subject on three occasions, separated by at least a 1 week wash-out period between each session. All subjects received both the tests

25 and reference drugs in a three-way crossover design.

Each subjects was randomly assigned at the first study period to either of the Test formulations or to the Reference formulation, and was subsequently crossed over at least one week later to either of the alternative treatments. The process was repeated during the third study session, such that each subject was exposed to one of the following

30 treatment schemes: $T_1 \rightarrow R \rightarrow T_2$; $T_1 \rightarrow T_2 \rightarrow R$; $R \rightarrow T_1 \rightarrow T_2$; $R \rightarrow T_2 \rightarrow T_1$; $T_2 \rightarrow R \rightarrow T_1$; $T_2 \rightarrow T_1 \rightarrow R$.

Drug concentration was determined by taking blood samples from all subjects regardless of treatment assignment at the following time points: 0 hour (pre-dosing), 0.5, 1, 1.5, 2, 3, 4, 6, 8, 10 and 12 hours post-initial dosing, for a total of 11 samples per study. Each sample was tested for simvastatin lactone and simvastatin hydroxyacid, the active metabolite, by analysis using a validated LC/MS/MS method.

The AUC_t and AUC_∞, C_{max}, T_{max}, and half life ($t_{1/2}$) were calculated for each volunteer both for simvastatin in plasma and for the active metabolite simvastatin hydroxyacid in plasma. Table 1 illustrates the average values for simvastatin in plasma and compares the values of the two test formulations to the average values obtained with the reference formulation.

Table 1: Simvastatin Concentration in Plasma					
	Test 1	Test 2	Reference	Test1/Ref.	Test2/Ref.
Avg. AUC _t (ng*h/ml)	20.8	26.9	14.9	1.4	1.81
Avg. AUC _∞ (ng*h/ml)	25.9	33.0	16.3	1.59	2.02
Avg. C _{max} (ng/ml)	5.8	8.0	5.6	1.04	1.43
Avg. T _{max} (hours)	1.58	2.46	1.04		
Avg. $t_{1/2}$ (hours)	3.99	3.10	2.03		

Table 1 demonstrates that both test formulations showed improved bioavailability over the simvastatin reference with the sample having 20 mg of simvastatin dissolved in 180 mg of menthol giving even better results than the concomitant dosing of a 20 mg simvastatin tablet along with a capsule of 180 mg of menthol. For AUC_t the average improvement in the bioavailability of Test 1 (concomitant separate dosing) the improvement was 40% while the improvement for the drug dissolved in menthol was 81%. The corresponding values for the AUC extrapolated to infinity were 59% and 102%, respectively. Consequently, the dissolved product gave larger improvements than concomitant separate dosing.

The ratio of the AUC_t of each test formulation to the reference formulation for each volunteer was calculated (each volunteer being his own control) and the average value of the ratio calculated. These results are illustrated in Table 2.

Table 2: Ratio Analysis of AUC _t for Simvastatin in Plasma					
Subject	Test 1	Test 2	Reference	Test 1/Ref.	Test 2/Ref.
1	20.78	21.43	27.75	0.749	0.772
2	29.54	39.70	38.96	0.758	1.02
3	23.53	17.62	6.76	3.48	2.61
4	26.89	75.45	33.12	0.812	2.28

5	37.32	15.08	6.68	5.59	2.26
6	12.7	10.85	4.31	2.95	2.52
7	8.18	6.26	5.42	1.51	1.16
8	13.48	26.46	13.85	0.975	1.91
9	31.96	66.15	13.53	2.36	4.89
10	22.29	23.63	15.29	1.46	1.55
11	14.66	9.55	7.75	1.89	1.23
12	8.79	10.68	5.17	1.70	2.07
Mean	20.8	26.9	14.9	2.02	2.02
± SD	9.4	22.5	11.9	1.42	1.09
CV%	45.3	83.7	80.1	70.5	54

Table 2 illustrates the ratio analysis of the AUCt values. Both test formulations showed a more than 100% improvement in bioavailability compared to the reference formulation. The two test formulations gave the same larger improvement. The value for Tmax is somewhat delayed for Test 2 compared to the reference and slightly so for Test 1. The values of $t_{1/2}$ are slightly longer, which may indicate competition by menthol for metabolic pathways that determine the $t_{1/2}$ such a glucuronidation and CYP3A4 pathways.

Table 3 collected the average values for simvastatin hydroxyacid, the active metabolite, in plasma and compared the values of the two test formulations to the average values obtained with the reference formulation.

Table 3: Simvastatin Hydroxyacid in Plasma					
	Test 1	Test 2	Reference	Test 1/Ref.	Test 2/Ref.
Avg AUCt (ng*h/ml)	10.2	12.7	8.9	1.15	1.43
Avg Cmax (ng/ml)	1.32	1.71	1.18	1.12	1.45
Avg Tmax (hours)	5.5	5.4	5.4		
Avg $t_{1/2}$ (hours)	8.8	5.7	6.5		

Table 3 illustrates the values for the active metabolite of simvastatin. Both test formulations showed improved bioavailability as expressed as average AUCt. Test 1 (concomitant separate dosing) showed a 15% improvement in the average bioavailability of the active moiety when compared to the reference drug product. Test 2 (concomitant dissolved dosing) showed a 45% improvement in the average AUCt and therefore in average bioavailability.

The ratio of AUCt of each test formulation to the reference formulation for each volunteer was calculated (each volunteer being his own control) and the average value of the ratio calculated. These results are illustrated in Table 4.

Table 4: Ratio Analysis of AUCt for Simvastatin Hydroxyacid in Plasma

Subject	Test 1	Test 2	Reference	Test 1/Ref.	Test 2/Ref.
1	9.75	7.44	10.49	0.927	0.710
2	17.28	19.10	25.99	0.665	0.735
3	9.57	11.66	5.20	1.84	2.24
4	7.85	18.49	8.04	0.976	2.30
5	13.23	9.05	5.52	2.40	1.64
6	7.61	11.3	5.18	1.47	2.18
7	9.55	10.88	7.29	1.31	1.49
8	3.76	5.46	4.18	0.900	1.31
9	9.65	17.77	4.70	2.06	3.78
10	19.24	21.77	16.37	1.18	1.22
11	12.69	13.61	10.97	1.16	1.24
12	2.64	6.37	2.73	0.967	2.33
Mean	10.2	12.7	8.9	1.32	1.77
± SD	4.9	5.4	6.6	0.53	0.85
CV%	47.5	42.5	73.9	39.8	48.1

Table 4 illustrates the ratio analysis for the AUCt values for the active moiety.

Both test formulations showed a clear improvement in the average of the individual ratios of AUCt with Test 2 being superior to Test 1. Test 1 showed an improved ratio of 32% compared to the reference drug product while Test 2 showed a 77% improvement. The variability of the drug absorption for the active moiety is also clearly improved when dosing with menthol. The reference had a percent coefficient of variation of 74% while Test 1 showed 48% and Test 2 43%, both a considerable improvement and Test 2 being superior.

Therefore, administering simvastatin with menthol can significantly improve the bioavailability of both the parent drug and its active metabolite and delivering the drug when dissolved in the menthol gives an even greater improvement in the improved bioavailability, and a lower variability of the active moiety. The approximate 80 to 100% improvement in the bioavailability of the simvastatin itself and the simvastatin hydroxyacid active moiety along with lowered variability should be able to lead to improved dosing and treatment with this important drug.

Claims

What is claimed is:

1. A composition for improving the bioavailability of a drug comprising at
5 least one poorly bioavailable drug dissolved in an effective amount of menthol.
2. The composition according to claim 1, wherein the poorly bioavailable
drug is a drug with low aqueous solubility, a drug metabolized by cytochrome P450, a
drug expelled from cells by the P-glycoprotein pump, or a drug metabolized via
10 glucuronidation.
3. The composition according to claim 2, wherein the drug with low aqueous
solubility is a drug having a water solubility of less than about 20 mg/ per milliliter of
water.
15
4. The composition according to claim 1, wherein the drug is cyclosporine,
atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pravastatin, simvastatin,
paclitaxel, fenofibrate, itraconazole, bromocriptine, carbamazepine, diazepam, paclitaxel,
etoposide, camptothecin, danazole, progesterone, nitrofurantoin, estradiol, estrone,
20 oxfendazole, proquazone, ketoprofen, nifedipine, verapamil, or glyburide.
5. The composition according to claim 1, wherein the drug is cyclosporine,
atorvastatin, cerivastatin, fluvastatin, lovastatin, mevastatin, pravastatin, simvastatin, or
paclitxel.
25
6. The composition according to claim 1, wherein the compound is
simvastatin, paclitaxel, or cyclosporine.
7. A method for improving the bioavailability of a drug comprising
30 dissolving the drug in an effective amount of menthol.
8. A method for improving the bioavailability of a drug comprising
dissolving at least one poorly bioavailable drug in an effective amount of menthol.

9. The method according to claim 8, wherein the poorly bioavailable drug is a drug with low aqueous solubility, a drug capable of being metabolized by cytochrome P450, a drug capable of being expelled from cells by the P-glycoprotein pump, or a drug capable of being metabolized via glucuronidation.

5

10. The method according to claim 8, further comprising administering the composition to a mammal.

11. The method according to claim 8, wherein the amount of menthol is sufficient to increase the oral bioavailability of the drug by an amount represented by an about 10% or more increase in the average area under the blood or plasma concentration versus time curve (AUC) when compared to a non-menthol containing formulation AUC.

12. The method according to claim 9, wherein the amount of menthol is about 60% to 99% by weight.

13. A method for reducing the variability of the bioavailability of a drug comprising dissolving at least one poorly bioavailable drug in an effective amount of menthol.

20

14. The method according to claim 13, wherein the poorly bioavailable drug is a drug with low aqueous solubility, a drug capable of being metabolized by cytochrome P450, a drug capable of being expelled from cells by the P-glycoprotein pump, or a drug capable of being metabolized via glucuronidation.

25

15. The method according to claim 13, further comprising administering the composition to a mammal.

16. The method according to claim 13, wherein the amount of menthol is sufficient to decrease the variability in the drug's bioavailability by about 10% or more of the relative standard deviation (CV%) of the area under the blood or plasma concentration versus time curve (AUC) when compared to a non-menthol containing formulation AUC.

30

17. A method for increasing the extent of time that a drug provides a therapeutically significant concentration in blood or plasma comprising dissolving at least one poorly bioavailable drug in an effective amount of menthol.

5 18. The method according to claim 17, wherein the poorly bioavailable drug is a drug with low aqueous solubility, a drug capable of being metabolized by cytochrome P450, a drug capable of being expelled from cells by the P-glycoprotein pump, or a drug capable of being metabolized via glucuronidation.

10 19. The method according to claim 17 wherein the amount of menthol is sufficient to extend the time that the drug provides a therapeutically significant concentration in blood or plasma by one hour or more.

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/073686 A3

(51) International Patent Classification⁷: **A61K 9/14**

(21) International Application Number:
PCT/US2004/004684

(22) International Filing Date: 17 February 2004 (17.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/449,246 20 February 2003 (20.02.2003) US

(71) Applicant (for all designated States except BB, US): **TEVA PHARMACEUTICAL INDUSTRIES LTD.** [IL/IL]; 5 Basel Street, P.O. Box 3190, 49131 Petah-Tiqva (IL).

(71) Applicant (for BB only): **TEVA PHARMACEUTICALS USA, INC.** [US/US]; 1090 Horsham Road, P.O. Box 1090, North Wales, PA 19454-1090 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FLASH-NER-BARAK, Moshe** [IL/IL]; Hefetz Mordechai 15, Petach Tikva (IL). **LERNER, Itzhak, E.** [IL/IL]; Wolfson 32, Petach Tiqva (IL). **ROSENBERGER, Vered** [IL/IL]; 3 Miss Landau Street, Jerusalem 96410 (IL). **MOLDAVSKI, Naomi** [IL/IL]; Moshav Dekel D.N., HaNegev 85492 (IL).

(74) Agent: **BRAINARD, Charles, R.**; Kenyon & Kenyon, One Broadway, New York, NY 10004-1050 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
4 November 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MENTHOL SOLUTIONS OF DRUGS

(57) Abstract: The present invention relates to compositions comprising solutions of drugs in menthol, especially drugs that are poorly soluble in water, and to methods for making such compositions.

WO 2004/073686 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/004684

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>NAZZAL S ET AL: "Preparation and in vitro characterization of a eutectic based semisolid self-nanoemulsified drug delivery system (SNEDDS) of ubiquinone: Mechanism and progress of emulsion formation"</p> <p>INTERNATIONAL JOURNAL OF PHARMACEUTICS (KIDLINGTON), vol. 235, no. 1-2, 20 March 2002 (2002-03-20), pages 247-265, XP002261220 ISSN: 0378-5173 abstract paragraph '2.2.1! paragraph '03.1!; figures 2,3</p> <p style="text-align: center;">----- -/-</p>	1-3,7-19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

31 August 2004

Date of mailing of the international search report

20/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Epskamp, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/004684

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STOTT P W ET AL: "Transdermal delivery from eutectic systems: enhanced permeation of a model drug, ibuprofen" JOURNAL OF CONTROLLED RELEASE, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, NL, vol. 50, no. 1-3, 2 January 1998 (1998-01-02), pages 297-308, XP004107662 ISSN: 0168-3659 abstract paragraph '03.1! paragraph '04.1!; figure 5 paragraph '04.3!; figure 10; table 1	1-3,7-19
X	KANG LISHENG ET AL: "Physicochemical studies of lidocaine-menthol binary systems for enhanced membrane transport" INTERNATIONAL JOURNAL OF PHARMACEUTICS (KIDLINGTON), vol. 206, no. 1-2, 25 September 2000 (2000-09-25), pages 35-42, XP002294480 ISSN: 0378-5173 abstract paragraph '02.2! paragraph '03.1!; figures 1,2 paragraph '03.5!; table 1	1-3,7-19
X	KAPLUN-FRISCHOFF Yael ET AL: "Testosterone skin permeation enhancement by menthol through formation of eutectic with drug and interaction with skin lipids" JOURNAL OF PHARMACEUTICAL SCIENCES, vol. 86, no. 12, December 1997 (1997-12), pages 1394-1399, XP001068689 ISSN: 0022-3549 abstract page 1394, right-hand column, paragraph 3 - paragraph 7 page 1395, right-hand column, last paragraph - left-hand column, paragraph 1; figures 1,2 page 1396, right-hand column, paragraph 2 - paragraph 3; figure 3	1-3,7-19
A	WO 96/40192 A (AVMAX INC ; BENET LESLIE Z (US); BENET REED M (US); WACHER VINCENT J ()) 19 December 1996 (1996-12-19) cited in the application page 2, last paragraph page 5, line 4 - page 6, last line ; tables 1,2 page 31, last paragraph claims	1-19

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/004684

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>WO 03/082247 A (ROSENBERGER VERED ; TEVA PHARMA (IL); DRABKIN ANNA (IL); LERNER ITZHAK) 9 October 2003 (2003-10-09) page 3, line 6 - line 16 examples 1-5 claims</p> <p>-----</p>	1-19

Form PCT/ISA/210 (continuation of second sheet) (January 2004)

BNSDOCID: <WO____2004073686A3_I_>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2004/004684

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 8-16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/004684

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640192	A	19-12-1996	US 5716928 A 10-02-1998
			US 5665386 A 09-09-1997
			CA 2224227 A1 19-12-1996
			EP 0831870 A1 01-04-1998
			JP 11507356 T 29-06-1999
			US 6121234 A 19-09-2000
			WO 9640192 A1 19-12-1996
			US 5916566 A 29-06-1999
WO 03082247	A	09-10-2003	WO 03082247 A2 09-10-2003
			US 2003224059 A1 04-12-2003

Form PCT/ISA/210 (patent family annex) (January 2004)